

Holdrege Veterinary Clinic

2201 4th AVE, Holdrege, NE 68949
308-995-6591

Canine Vaccination Protocol Puppy 6 to 16 weeks old

If under 8 weeks of age at 1st visit

DAPPCV (then not needed at 20 week visit)
Deworm

8 weeks or 1st visit

DAPPCV*
Bordatella (Kennel Cough)
Deworm
Flea/Tick Control (Frontline Plus)**

12 weeks or 2nd visit

DAPPCV
Bordatella Booster
Rabies
Deworm
Flea/Tick Control
Heartworm prevention***

16 weeks or 3rd visit

DAPPCV
Lepto Vaccine****

20 weeks or 4th visit

Lepto Vaccine Booster
DAPPCV

4-6 months

Spay or neuter

Yearly Boosters

DAPPV
Rabies (booster every 2 years after the first two vaccinations)
Bordatella
Heartworm test
4 weeks later: Lepto Booster

*DAPPCV (Distemper, Adenovirus, Parvovirus, Parainfluenza, and Corona Virus)

**Flea and Tick Control/Prevention should be administered monthly during flea and tick season

***Heartworm Prevention should be administered all year round

****Leptosporosis is a bacteria that is shed in the urine of wildlife and rodents. Working dogs (hunting, cattle, and on farm) have a high risk of exposure and can become infected by ingesting contaminated water in ponds and streams. Lepto can cause liver failure, kidney failure, or both. It can be extremely expensive and difficult to treat. Lepto can also infect humans.

The booster needs to be given within 4 weeks of the initial vaccination for optimal response.

An **Annual Wellness Exam** with fecal and urinalysis is recommended with your annual vaccinations.

An **Annual Geriatric Blood Profile Test** is recommended for patients 7 years of age and older for early detection of diabetes, kidney, and liver disease and many other health problems.