

Holdrege Veterinary Clinic
2201 4th AVE, Holdrege, NE 68949
308-995-6591

Canine Vaccination Protocol

**Puppy over 16 weeks of age or
adult with unknown vaccine history**

1st Appointment

DAPPV*
Bordatella
Rabies
Deworm
Flea/Tick Control**
Heartworm test and prevention ***

2nd Appointment (3-4 weeks later)

DAPPV
Deworm
Lepto Vaccine****
Bordatella Booster

3rd Appointment (3-4 weeks later)

DAPPV
Lepto Vaccine Booster

Yearly Boosters

DAPPV
Rabies (boosted every 2 years after the first two vaccinations)
Bordatella
Heartworm test
4 weeks later: Lepto Booster

*DAPPCV (Distemper, Adenovirus, Parvovirus, Parainfluenza, and Corona Virus)

**Flea and Tick Control/Prevention should be administered monthly during flea and tick season

***Heartworm Prevention should be administered all year round

****Leptosporosis is a bacteria that is shed in the urine of wildlife and rodents. Working dogs (hunting, cattle, and on farm) have a high risk of exposure and can become infected by ingesting contaminated water in ponds and streams. Lepto can cause liver failure or kidney failure, or both. It can be extremely expensive and difficult to treat. Lepto can also infect humans.

The booster needs to be given within 4 weeks of the initial vaccination for optimal response.

An **Annual Wellness Exam** with fecal and urinalysis is recommended with your annual vaccinations.

An **Annual Geriatric Blood Profile Test** is recommended for patients 7 years of age and older for early detection of diabetes, kidney, and liver disease and many other health problems.